

Hasil Analisis Statistik

Case Processing Summary

	Valid		Cases Missing		Total	
	N	Percent	N	Percent	N	Percent
Kebiasaan Mencuci Tangan * Kontaminasi STH	46	100.0%	0	0.0%	46	100.0%

Kebiasaan Mencuci Tangan * Kontaminasi STH Crosstabulation

			Kontaminasi STH		Total
			Negatif	Positif	
Kebiasaan Mencuci Tangan	Ya	Count	40	4	44
		% within Kebiasaan Mencuci Tangan	90.9%	9.1%	100.0%
	Tidak	Count	2	0	2
		% within Kebiasaan Mencuci Tangan	100.0%	0.0%	100.0%
Total	Count		42	4	46
	% within Kebiasaan Mencuci Tangan		91.3%	8.7%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.199 ^a	1	.655		
Continuity Correction ^b	.000	1	1.000		
Likelihood Ratio	.372	1	.542		
Fisher's Exact Test				1.000	.832
Linear-by-Linear Association	.195	1	.659		
N of Valid Cases	46				

a. 3 cells (75.0%) have expected count less than 5. The minimum expected count is .17.

b. Computed only for a 2x2 table

Case Processing Summary

	Valid		Cases Missing		Total	
	N	Percent	N	Percent	N	Percent
Kebiasaan Lokasi Defekasi * Kontaminasi STH	46	100.0%	0	0.0%	46	100.0%

Kebiasaan Lokasi Defekasi * Kontaminasi STH Crosstabulation

			Kontaminasi STH		Total
			Negatif	Positif	
Kebiasaan Lokasi Defekasi	Jamban	Count	36	4	40
		% within Kebiasaan Lokasi Defekasi	90.0%	10.0%	100.0%
	Sungai/Tanah	Count	6	0	6
		% within Kebiasaan Lokasi Defekasi	100.0%	0.0%	100.0%
Total	Count		42	4	46
	% within Kebiasaan Lokasi Defekasi		91.3%	8.7%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.657 ^a	1	.418		
Continuity Correction ^b	.001	1	.973		
Likelihood Ratio	1.174	1	.279		
Fisher's Exact Test				1.000	.560
Linear-by-Linear Association	.643	1	.423		
N of Valid Cases	46				

a. 2 cells (50.0%) have expected count less than 5. The minimum expected count is .52.

b. Computed only for a 2x2 table

Case Processing Summary

	Valid		Cases Missing		Total	
	N	Percent	N	Percent	N	Percent
Kebiasaan Memotong Kuku * Kontaminasi STH	46	100.0%	0	0.0%	46	100.0%

Kebiasaan Memotong Kuku * Kontaminasi STH Crosstabulation

			Kontaminasi STH		Total
			Negatif	Positif	
Kebiasaan Memotong Kuku	Sering	Count	27	3	30
		% within Kebiasaan Memotong Kuku	90.0%	10.0%	100.0%
	Jarang	Count	15	1	16
		% within Kebiasaan Memotong Kuku	93.8%	6.3%	100.0%
Total	Count		42	4	46
	% within Kebiasaan Memotong Kuku		91.3%	8.7%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.185 ^a	1	.667		
Continuity Correction ^b	.000	1	1.000		
Likelihood Ratio	.194	1	.660		
Fisher's Exact Test				1.000	.566
Linear-by-Linear Association	.181	1	.671		
N of Valid Cases	46				

a. 2 cells (50.0%) have expected count less than 5. The minimum expected count is 1.39.

b. Computed only for a 2x2 table

Case Processing Summary

	Valid		Cases Missing		Total	
	N	Percent	N	Percent	N	Percent
Kepemilikan Jamban * Kontaminasi STH	46	100.0%	0	0.0%	46	100.0%

Kepemilikan Jamban * Kontaminasi STH Crosstabulation

			Kontaminasi STH		Total
			Negatif	Positif	
Kepemilikan Jamban	Ya	Count	38	4	42
		% within Kepemilikan Jamban	90.5%	9.5%	100.0%
	Tidak	Count	4	0	4
		% within Kepemilikan Jamban	100.0%	0.0%	100.0%
Total	Count		42	4	46
	% within Kepemilikan Jamban		91.3%	8.7%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.417 ^a	1	.518		
Continuity Correction ^b	.000	1	1.000		
Likelihood Ratio	.763	1	.382		
Fisher's Exact Test				1.000	.686
Linear-by-Linear Association	.408	1	.523		
N of Valid Cases	46				

a. 3 cells (75.0%) have expected count less than 5. The minimum expected count is .35.

b. Computed only for a 2x2 table

Case Processing Summary

	Valid		Cases Missing		Total	
	N	Percent	N	Percent	N	Percent
Jenis Pembuangan Tinja * Kontaminasi STH	46	100.0%	0	0.0%	46	100.0%

Jenis Pembuangan Tinja * Kontaminasi STH Crosstabulation

			Kontaminasi STH		Total
			Negatif	Positif	
Jenis Pembuangan Tinja	Septic tank	Count	34	1	35
		% within Jenis Pembuangan Tinja	97.1%	2.9%	100.0%
	Sungai/Selokan	Count	8	3	11
		% within Jenis Pembuangan Tinja	72.7%	27.3%	100.0%
Total	Count		42	4	46
	% within Jenis Pembuangan Tinja		91.3%	8.7%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	6.284 ^a	1	.012		
Continuity Correction ^b	3.585	1	.058		
Likelihood Ratio	5.208	1	.022		
Fisher's Exact Test				.037	.037
Linear-by-Linear Association	6.147	1	.013		
N of Valid Cases	46				

a. 2 cells (.50.0%) have expected count less than 5. The minimum expected count is .96.

b. Computed only for a 2x2 table

Mantel-Haenszel Common Odds Ratio Estimate

Estimate		12.750
ln(Estimate)		2.546
Standard Error of ln(Estimate)		1.220
Asymptotic Significance (2-sided)		.037
Asymptotic 95% Confidence Interval	Common Odds Ratio	Lower Bound 1.168
		Upper Bound 139.235
	ln(Common Odds Ratio)	Lower Bound .155
		Upper Bound 4.936

The Mantel-Haenszel common odds ratio estimate is asymptotically normally distributed under the common odds ratio of 1.000 assumption. So is the natural log of the estimate.

Case Processing Summary

	Valid		Cases Missing		Total	
	N	Percent	N	Percent	N	Percent
Jenis Sumber Air * Kontaminasi STH	46	100.0%	0	0.0%	46	100.0%

Jenis Sumber Air * Kontaminasi STH Crosstabulation

			Kontaminasi STH		Total
			Negatif	Positif	
Jenis Sumber Air	PDAM/Sumur bor	Count	10	0	10
		% within Jenis Sumber Air	100.0%	0.0%	100.0%
	Sumur gali	Count	32	4	36
		% within Jenis Sumber Air	88.9%	11.1%	100.0%
Total	Count		42	4	46
	% within Jenis Sumber Air		91.3%	8.7%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1.217 ^a	1	.270		
Continuity Correction ^b	.220	1	.639		
Likelihood Ratio	2.064	1	.151		
Fisher's Exact Test				.562	.361
Linear-by-Linear Association	1.190	1	.275		
N of Valid Cases	46				

a. 2 cells (50.0%) have expected count less than 5. The minimum expected count is .87.

b. Computed only for a 2x2 table

Case Processing Summary

	Valid		Cases Missing		Total	
	N	Percent	N	Percent	N	Percent
Jarak Pembuangan Tinja > 10 m * Kontaminasi STH	46	100.0%	0	0.0%	46	100.0%

Jarak Pembuangan Tinja > 10 m * Kontaminasi STH Crosstabulation

			Kontaminasi STH		Total
			Negatif	Positif	
Jarak Pembuangan Tinja > 10 m	Ya	Count	23	2	25
		% within Jarak Pembuangan Tinja > 10 m	92.0%	8.0%	100.0%
	Tidak	Count	19	2	21
		% within Jarak Pembuangan Tinja > 10 m	90.5%	9.5%	100.0%
Total	Count		42	4	46
	% within Jarak Pembuangan Tinja > 10 m		91.3%	8.7%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.033 ^a	1	.855		
Continuity Correction ^b	.000	1	1.000		
Likelihood Ratio	.033	1	.855		
Fisher's Exact Test				1.000	.626
Linear-by-Linear Association	.033	1	.857		
N of Valid Cases	46				

a. 2 cells (50.0%) have expected count less than 5. The minimum expected count is 1.83.

b. Computed only for a 2x2 table