

## Neglected Posterior Subtalar Dislocation: A Case Report and Operative Technique

Aditya Priherdadi, Raden Andri Primadhi

Department of Orthopaedics dan Traumatology Faculty of Medicine Universitas Padjadjaran

Dr. Hasan Sadikin General Hospital Bandung, Indonesia

### Abstract

Posterior subtalar dislocation is an exceedingly rare entity. Early anatomic reduction, initially by closed means, is crucial. Failure to recognize this condition or any occurrence of neglect will complicate the management and worsen the outcome. There are a handful of case reports on posterior subtalar dislocation and even less for neglected cases. This study reports a neglected subtalar dislocation in a 26-year-old male who worked as a driver and had experienced high-energy trauma due to car collision. This patient was managed by open reduction and pinning, and was followed up 2 months later with a good outcome. This technique was selected to be able to achieve functional motions without pain.

**Keywords:** complication, intractable pain, open reduction, subtalar dislocation.

## Dislokasi Posterior Sendi Subtalar yang Terlantar: Laporan Kasus dan Teknik Operasi

### Abstrak

Dislokasi posterior sendi subtalar adalah hal yang sangat jarang terjadi. Reduksi secara anatomis pada awal kejadian, secara tertutup, sangat krusial. Kegagalan untuk mengenali kondisi ini atau tidak tepat penanganan akan menyulitkan manajemen dan memperburuk hasilnya. Hanya ada sedikit laporan kasus tentang dislokasi posterior sendi subtalar dan untuk kasus yang terlantar lebih jarang lagi ditemukan. Penelitian ini melaporkan sebuah kasus dislokasi posterior sendi subtalar yang terlantar pada pria berusia 26 tahun yang bekerja sebagai pengemudi dan mengalami trauma karena tabrakan mobil. Dilakukan reduksi terbuka dan pemasangan pin, serta ditindaklanjuti 2 bulan kemudian dengan hasil yang memuaskan. Teknik ini bertujuan untuk mencapai gerakan sendi yang fungsional tanpa nyeri.

**Kata kunci:** Komplikasi, nyeri berat, reduksi terbuka, dislokasi subtalar

---

**Corresponding Author:** Aditya Priherdadi, Department of Orthopaedics dan Traumatology Faculty of Medicine Universitas Padjadjaran/Dr. Hasan Sadikin General Hospital Bandung, Jalan Pasteur No. 38 Bandung, West Java, Indonesia, Email: priherdadi@gmail.com

## Introduction

Subtalar joint dislocation is an uncommon injury, and cases of posterior subtalar dislocation are even more rare. This injury is characterized by a simultaneous dislocation of talocalcaneal and talonavicular joints, while the tibiotalar and calcaneocuboid articulations remains intact.<sup>1</sup> Under-diagnosis of this type of cases by primary health care providers may occur due to the rarity of this injury. Delayed diagnosis will delay proper treatment. Unsuccessful attempt of reduction could eventually result in severe swelling, intractable pain, vascular problems, and damages to the articular surfaces. This case highlights the clinical and imaging features of posterior subtalar dislocation, complications from neglect, reduction techniques with Schanz screw, and fixation with Kirschner wire that providers should be familiar with.

## Case

A 26 years-old male minibus driver sustained an injury to his foot in a car collision one week prior to hospital admission. The exact mechanism was unknown, possibly a fastened foot during the forward flip of the body. He was brought to the nearest hospital, underwent radiological

examinations, and was diagnosed to experience posterior subtalar dislocation. An unsuccessful attempt of closed reduction was performed. The doctor then suggested him to go to the referral hospital, but the patient refused the medical advice and went home to seek help from a traditional bone setter instead.

One week later, he was presented to our hospital due to the intractable pain and severe swelling. Physical examination revealed a slight plantarflexion without neurovascular compromise (Figure 1). There were sores at the medial side, possibly from the initial trauma or the attempted reduction. He had no prior injuries or surgery to this ankle.

Plain radiographs and sagittal CT Scan of the foot revealed a posterior subtalar dislocation with the calcaneocuboid joint remained intact (Figure 2a, 2b). The axial CT Scan showed a gap between talar body and calcaneus without any horizontal translation (Figure 2c). A displaced trigonum bone posterior to the talus was also observed.

On physical examinations, the subtalar dislocation was difficult to observe because there was no obvious medial or lateral translation of the calcaneus. Among the first differential diagnoses were fractures around foot and ankle, severe ankle sprain, or soft tissue contusions.

Imaging studies showed a posterior subtalar



Figure 1 Clinical Picture Upon Arrival




**Figure 2** Imaging Studies (2a: Plain X-ray, 2b: Sagittal CT scan, 2c: Axial CT scan)

dislocation. Considering the acute setting, severe swelling, impinged tissue, and buttonholing bone were listed as possible reduction obstacles.

It was decided that the patient should undergo surgical reduction. Surgical indication for this case was the unsuccessful closed reduction

attempt. Considering the swollen foot, it was impossible to do another attempt.

A Schanz screw was inserted to the calcaneus as a joystick for reduction. The calcaneus was pulled downward and anteriorly to facilitate reduction (Figure 3). The talar head was



**Figure 3** Intraoperative Reduction



**Figure 4 Immediate Post-operative X-ray**

buttonholing to the surrounding tissue and hindered the reduction. An incision was made at the talonavicular joint to release the soft tissue. Hyperplantar flexion aims to reduce the strain on the flexor bundle and achilles so that the calcaneus is easy to move and gives the talonavicular joint distance so that the head of the talus is easily reduced.

After the subtalar and talonavicular joints were restored, several Kirschner wires was put across the posterior facet for temporary fixation (Figure 4). The patient was sent home three days post-surgery after the pain subsided and there was no sign of infection.

The wound healed uneventfully. At the 1-month follow-up the Kirschner wires were removed and the patient was able to walk in partial weight bearing painlessly. Full weight bearing was achieved as advised at the 2-months follow-up. The functional results of the patient was evaluated according to Gay and Evrard's criteria based on a study of pain, walking, mobility, stability, and professional activities. Very good functional results with pain 3, instability 3, mobility 2, edema 2, occupation 3, leading to an accumulative 13 points were observed.

## Discussion

Subtalar dislocation is uncommon and accounts

for approximately 1% of all dislocations. The talonavicular and talocalcaneal joints are involved simultaneously, while the tibiotalar and calcaneocuboid joints remain intact.<sup>2,3,4</sup> There is a strong association between subtalar dislocation and high energy trauma, such as a fall from height or road traffic accidents. The mechanism of posterior dislocation is hyperplantarflexion of the foot. Both medial and lateral ligaments, along with the interosseus ligament, are torn in posterior and anterior dislocations.<sup>5,6,7</sup> This patient's foot was slightly plantarflexed, without any locked supination or pronation. No rotational component was found in the calcaneus in this patient. This findings suggested that the trauma was in pure hyperplantar flexion.<sup>1</sup>

In this case, talar head buttonholing is one among the obstacles for reduction, other than the fibrosis and swollen condition the foot. Similar to the irreducible medial subtalar dislocation, the talar head buttonholed through the extensor digitorum brevis (EDB) muscles and extensor retinaculum. There was also an EDB muscle interpositioning between talus and navicular.<sup>8</sup> The chance of repositioning the joint by closed manipulation diminished considerably over time, while complications associated with closed reduction substantially increased due to the presence of fibrotic soft tissues.<sup>9</sup>

Following an unsuccessful attempt of closed reduction, open reduction is the next step. Release of capsule and other soft tissue is

necessary to reduce the joint without tension. For this patient, an incision at talonavicular joint was performed to release the buttonholing talar head and EDB tendon interposition. The Schanz screw inserted to the calcaneus was beneficial to facilitate reduction as a joystick while avoiding further soft tissue injuries as if the reduction was done by grasping the calcaneus. Talonavicular joint reduction was done simultaneously by simple manipulation using an elevator. After a stable reduction, an immobilisation was applied for 3-6 weeks using an external support or internal fixation. The duration of immobilization varies as there is a balance between avoiding instability and avoiding stiffness.<sup>10</sup> For this patient, it was decided to place several Kirschner wires to temporarily fix the subtalar joint. The posterior cast splint as an external support was another option for this condition; however, it was opted out in this case to facilitate wound care and accommodate swelling.

The goals for treating this dislocation is to prevent re-dislocation, recovery of an acceptable range of motion and strength of the extremity, in order to at least maintain daily activity.<sup>11</sup>

Complications for this condition include avascular necrosis of the talus, infection, posttraumatic arthritis, chronic subtalar instability, and complex regional pain syndrome with delayed reduction. Negative prognostic factors include lateral and open dislocations, total talar dislocations, and associated fractures.<sup>13</sup> A CT-scan is advisable after reduction to document additional bony injury to the subtalar area and adjacent bones.<sup>12</sup>

In conclusion, the posterior subtalar dislocation is very rare and the mild deformity in this condition may result in under-diagnosis by healthcare providers and patient's reluctance to undergo treatment. Unsuccessful attempts of closed reduction should prompt an investigation of the reduction obstacles and subsequent management, including surgical open reduction. Investigating talar head buttonholing to surrounding tissue is important. Tools such as Schanz screw are beneficial to facilitate reduction without further compromising soft tissues. Combining this technique with pinning K-wire fixation is a simple technique that can be performed by health care providers with good results.

## References

1. Thomas PC. Subtalar dislocation case report

- and literature review. *Global J Med Res.* 2016;2(1):19–23.
2. Jungbluth P, Wild M, Hakimi M, et al. Isolated subtalar dislocation. *J Bone Joint Surg Am.* 2010;92(4):890–4.
  3. Krähenbühl N, Horn-Lang T, Hintermann B, Knupp M. The subtalar joint: A complex mechanism. *EFORT Open Rev.* 2017;2(7):309–16.
  4. Yglesias B, Andrews K, Hamilton R, Lea J, Shah R, Ebraheim N. Case report: irreducible medial subtalar dislocation with incarcerated anterior talar head fracture in a young patient. *J Surg Case Rep.* 2018;2018(7):rjy168.
  5. Gaba S, Kumar A, Trikha V, Das S, Agrawal P. Posterior dislocation of subtalar joint without associated fracture: a case report and review of literature. *J Clin Diagn Res.* 2017;11(9):RD01–2.
  6. Sahil G, Arvind K, Vivek T, Saubhik D, Prabhat A. Posterior dislocation of subtalar joint without associated fracture: a case report and review of literature. *Journal of Clinical and Diagnostic Research.* 2017;11(9):1–2.
  7. Bali K, Kumar V, Bhagwat K, Rawall S. Closed posterior subtalar dislocation without any associated fracture: a case report and review of the literature. *Foot Ankle Surg.* 2011;17(3):e40–2.
  8. Azarkane M, Boussakri H, Alayyoubi A, Bachiri M, Elibrahimi A, Elmrini A. Closed medial total subtalar joint dislocation without ankle fracture: a case report. *J Med Case Rep.* 2014;8:313.
  9. Chung H, Yoon YS, Shin JS, Shin JJ, Kim D. Chronic irreducible anterior dislocation of the shoulder without significant functional deficit. *Clin Orthop Surg.* 2016;8(3):333–8.
  10. Mechchat A, Abdelhak M, Shimi M, Elibrahimi A, Elmrini A. Bilateral subtalar dislocation in gymnast: a case report. *Open Journal of Orthopedics.* 2013;3:258–60.
  11. Rammelt S, Goronzy J. Subtalar dislocations. *foot ankle Clin N Am.* 2015;20(2):253–64.
  12. Veltman ES, Steller EJA, Wittich P, Keizer J. Lateral subtalar dislocation: case report and review of the literature. *World J Orthop.* 2016;7(9):623–7.
  13. Prada-Cañizares A, Auñón-Martín I, Vilá Y Rico J, Pretell-Mazzini J. Subtalar dislocation: management and prognosis for an uncommon orthopaedic condition. *Int Orthop.* 2016;40(5):999–1007.